

The following samples were collected from Williamson Hall rooms:

- 311D
- 3<sup>rd</sup> floor main stairwell



Report for:

**Mr. Chad Johnson**  
**Eastern Washington University**  
EH&S, 002 Martin Hall  
Cheney, WA 99004

---

Regarding: Project: WIL  
EML ID: 1933913

Approved by:

Operations Manager  
Joshua Cox

Dates of Analysis:  
Direct microscopic exam (Qualitative): 05-30-2018

Service SOPs: Direct microscopic exam (Qualitative) (EM-MY-S-1039)  
AIHA-LAP, LLC accredited service, Lab ID #102297

---

All samples were received in acceptable condition unless noted in the Report Comments portion in the body of the report. Due to the nature of the analyses performed, field blank correction of results is not applied. The results relate only to the items tested.

EMLab P&K ("the Company") shall have no liability to the client or the client's customer with respect to decisions or recommendations made, actions taken or courses of conduct implemented by either the client or the client's customer as a result of or based upon the Test Results. In no event shall the Company be liable to the client with respect to the Test Results except for the Company's own willful misconduct or gross negligence nor shall the Company be liable for incidental or consequential damages or lost profits or revenues to the fullest extent such liability may be disclaimed by law, even if the Company has been advised of the possibility of such damages, lost profits or lost revenues. In no event shall the Company's liability with respect to the Test Results exceed the amount paid to the Company by the client therefor.

EMLab P&K's LabServe® reporting system includes automated fail-safes to ensure that all AIHA-LAP, LLC quality requirements are met and notifications are added to reports when any quality steps remain pending.

Client: Eastern Washington University  
 C/O: Mr. Chad Johnson  
 Re: WIL

Date of Sampling: 05-24-2018  
 Date of Receipt: 05-25-2018  
 Date of Report: 05-30-2018

**DIRECT MICROSCOPIC EXAMINATION REPORT**

Background Debris and/or Description	Miscellaneous Spores Present*	MOLD GROWTH: Molds seen with underlying mycelial and/or sporulating structures†	Other Comments††	General Impression
Lab ID-Version‡: 9101177-1, Analysis Date: 05/30/2018: Tape sample 1: WIL 311D				
Moderate	Very few	None	None	Normal trapping
Lab ID-Version: 9101178-1, Analysis Date: 05/30/2018: Tape sample 2: WIL 3rd Fl St Well W				
Moderate	Very few	None	None	Normal trapping

\* Indicative of normal conditions, i.e. seen on surfaces everywhere. Includes basidiospores (mushroom spores), myxomycetes, plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating. Distribution of spore types seen mirrors that usually seen outdoors.

† Quantities of molds seen growing are listed in the MOLD GROWTH column and are graded <1+ to 4+, with 4+ denoting the highest numbers.

†† Some comments may refer to the following: Most surfaces collect a mix of spores which are normally present in the outdoor environment. At times it is possible to note a skewing of the distribution of spore types, and also to note "marker" genera which may indicate indoor mold growth. Marker genera are those spore types which are present normally in very small numbers, but which multiply indoors when conditions are favorable for growth.

‡ A "Version" indicated by -"x" after the Lab ID# with a value greater than 1 indicates a sample with amended data. The revision number is reflected by the value of "x".  
 The limit of detection is < 1+ when mold growth is detected.

Client: Eastern Washington University  
 C/O: Mr. Chad Johnson  
 Re: WIL

Date of Sampling: 05-24-2018  
 Date of Receipt: 05-25-2018  
 Date of Report: 05-30-2018

**Mold/Fungal Growth Rating Details**

Growth Rating	Quantities of molds indicating growth are listed in the MOLD/FUNGAL GROWTH section. Judgement is used in determining the amount of growth present in the sample. For example, if only one portion of the sample has evidence of heavy growth, then it will receive a rating of heavy growth even though, strictly speaking, on a percentage basis of the entire sample, the amount of growth is low.	
	Swab/Tape/Dust/Wipe sample	Bulk Sample
< 1+ (Very Light Growth)	Evidence of very light growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in less than 10% of the microscopic fields examined.	Areas of very light growth detected by the presence of spores of one type seen with underlying mycelial and/or with their sporulating structures in the bulk sample.
1+ (Light Growth)	Evidence of light growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in 10 to 25% of the microscopic fields examined.	Areas of light growth detected by the presence of spores of one type seen with underlying mycelial and/or with their sporulating structures in the bulk sample.
2+ (Moderate Growth)	Evidence of moderate growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in 26 to 50% of the microscopic fields examined.	Areas of moderate growth detected by the presence of spores of one type seen with underlying mycelial and/or with their sporulating structures in the bulk sample.
3+ (Heavy Growth)	Evidence of heavy growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found in 51 to 75% of the microscopic fields examined.	Areas of heavy growth detected by the presence of spores of one type seen with underlying mycelial and/or with their sporulating structures in the bulk sample.
4+ (Very Heavy Growth)	Evidence of very heavy growth observed on the sample as indicated by spores of one type seen with underlying mycelial and/or with their sporulating structures found to be nearly confluent in the majority of the microscopic fields examined.	Areas of very heavy growth detected by the presence of spores of one type seen with underlying mycelial and/or with their sporulating structures in the bulk sample.

**Miscellaneous Spores**

Slides/specimens are examined for the presence of mold spores and pollen, noting the quantities and distribution of spore types found. A designation of 'normal trapping' is made when a mix of spore types is present with the same general distribution as is usually found outdoors. In other words, the biological component of the sample surface is like that found everywhere. Types of spores present would include basidiospores (mushroom spores), myxomycetes (slime molds), plant pathogens such as ascospores, rusts and smuts, and a mix of saprophytic genera with no particular spore type predominating. Many of these spore types would not be found growing indoors on building materials since many plant pathogens require living plants for growth, and mushrooms require compost, leaf duff of various types, or associations with roots of certain trees, etc. Due to these factors, when a mix of spores seen include these types as well as pollen, the rational source is the outside air, rather than indoor mold growth. The numbers of miscellaneous spores seen are graded and described as shown below as none, very few, few, variety, and wide variety.

None	Very Few	Few	Variety	Wide Variety
No spores detected	Very few spores detected	A few spores detected	Many spores containing a variety of different genera detected	Many spores containing a wide variety of different genera detected